

## “Convolution of an integral equation with the H-function as its kernel”

Ram Niwas Meghwal

Department of mathematics Govt. College Sujangarh, Rajasthan (India).

**Abstract** The object of this paper is to solve an integral equation of convolution form having H-function of two variable as it's kernel. Some known results are obtained as special cases

**Keywords** - integral equation, convolution, generalized type geometric function of two variables

**AMS Subject classification No.** 45E10

### 1. Definition and Introduction

The following definition and results will be required in this paper

(i) The Laplace Transform if

$$F(P) = L[f(t); p] = \int_0^{\infty} e^{-pt} f(t) dt, \quad \text{Re}(p) > 0 \quad \dots(1.1)$$

1.1)

Then F(p) is called the Laplace transform of f(t) with parameter p and is represented by  $F(p) = \int_0^{\infty} e^{-pt} f(t) dt$  Erdelyi [(3) pp.129-131 ]

(ii)  $L[f(t); p] = F(P)$  then  $L[e^{-at} f(t)] = F(p + a)$  ...

1.2)

And if  $f(0) = f'(0) = f''(0) = \dots = f^{m-1}(0) = 0$ ,  $f^n(t)$

Is continuous and differential, then

$$L[f^n(t); p] = P^n F(p)$$

....(1.3)

(iii) If  $L[f_1(t)] = F_1(p)$  then  $L[f_2(t)] = F_2(p)$

Then convolution theorem for Laplace transform is

$$L\left\{\int_0^1 f_1(t) f_2(t-u) du\right\} = L\{f_1(t)\} L\{f_2(t)\} = F_1(p) \cdot F_2(p)$$

... (1.4)

(iv) The H-Function Defined by Saxena and kumbhat [1] is an extension of Fox's H-Function on specializing the parameters, H-Function can be reduced to almost all the known special function as well as unknown

The Fox's H-Function of one variable is defined and represented in this Paper as follows [see Srivastava et al [2], pp 11-13 ]

$$H[x] = H_{P,Q}^{M,N} \left[ x / \begin{matrix} (a_j, \alpha_j)_{1, P} \\ (b_j, \beta_j)_{1, Q} \end{matrix} \right] = \frac{1}{2\pi\omega} \int_{\theta=N-1} \theta(\xi) x^\xi d\xi \quad \dots(1.5)$$

$$\theta(\xi) = \frac{\prod_{i=1}^n \Gamma b_j - \beta_j \xi \prod_{j=1}^N \Gamma 1 - a_j - \alpha_j \xi}{\prod_{i=M+1}^Q \Gamma 1 - b_j + \beta_j \xi \prod_{j=N+1}^P \Gamma a_j - \alpha_j \xi} \quad \dots(1.6)$$

For condition of the H-Function of one variable (1.5) and on the contour L we refer to srivastava et al [2] (V) The H-Function of two variable occurring in this paper is defined and represented as follows [see Srivastava et al [2] ,pp 83-85 ]

$$H[x, y] = H_{p_1, q_1, p_2, q_2, p_3, q_3}^{0, n_1, m_2, n_2, m_3, n_3} \left[ x y / \begin{matrix} (a_j, \alpha_j, A_j)_{1, p_1} & (c_j, z_j)_{1, p_2} & (e_j, E_j)_{1, p_3} \\ (b_j, \beta_j, B_j)_{1, q_1} & (d_j, \delta_j)_{1, q_2} & (f_j, F_j)_{1, q_3} \end{matrix} \right] \quad \dots(1.7)$$

$$= -\frac{1}{4\pi^2} \int_{L_1} \int_{L_2} \phi_1(\xi, \eta) \psi_2(\xi) \psi_3(\eta) x(\xi) y(\eta) d\xi d\eta \quad \text{Where}$$

$$\phi_1(\xi, \eta) = \prod_{j=1}^{n_1} \Gamma(1 - a_j + \alpha_j \xi + A_j \eta) \times \left[ \prod_{j=n_1+1}^{p_1} \Gamma a_j - \alpha_j \xi - A_j \eta \prod_{j=1}^{q_1} \Gamma 1 - b_j - \beta_j \xi + B_j \eta \right] \quad \dots(1.8)$$

Where the  $\psi_2(\xi)$  and  $\psi_3(\eta)$  are defined as (1.6) and for conditions of existence etc. of the  $H(x, y)$  we refer to srivastava et al [2]

### 2. Main Result

**Result I**  $L \left\{ t^\alpha H_{P,Q}^{M,N} \left[ at^\alpha / \begin{matrix} (a_j, \alpha_j)_{1, P} \\ (b_j, \beta_j)_{1, Q} \end{matrix} \right], P \right\}$   
 $= P^{-1-\alpha} H_{P+1, Q}^{M, N+1} \left[ at^{-\lambda} / \begin{matrix} (-\alpha, \lambda) (a_j, \alpha_j)_{1, P} \\ (b_j, \beta_j)_{1, Q} \end{matrix} \right]$

Provided  $\text{Re}(p) > 0 \ 1 \geq \lambda > a$  and  $\text{Re}(1+\alpha) > 0$

**Result II**

$$\int_0^1 \left\{ x^{\alpha-1} (1-x)^{\beta-1} H_{P_1, Q_1}^{M_1, N_1} \left[ z_1 x^\lambda / \begin{matrix} (a_j, \alpha_j)_{1, P_1} \\ (b_j, \beta_j)_{1, Q_1} \end{matrix} \right] H_{P_2, Q_2}^{M_2, N_2} \left[ z_2 (1-x)^\mu / \begin{matrix} (c_j, \gamma_j)_{1, P_2} \\ (d_j, \delta_j)_{1, Q_2} \end{matrix} \right] \right\} dx$$

$$= H_{P+2, Q+1}^{0, N+2, M_1, N_1, M_2, N_2} \left[ \begin{matrix} z_1 / (1-\alpha, \lambda) (1-\beta, \mu) (a_j, \alpha_j)_{P_1} (c_j, \gamma_j)_{P_2} \\ z_2 / (1-b-\beta, \lambda; \mu) (b_j, \beta_j)_{Q_1} (d_j, \delta_j)_{Q_2} \end{matrix} \right]$$

Provided  $\text{Re}(\alpha) > 0 \ \text{Re}(\beta) > 0 \ \lambda, \mu > 0$

$$\text{Re} \left( \alpha + \lambda \frac{b_j}{\beta_j} \right) > 0 \ \text{Re} \left( \beta + \mu \frac{d_j}{\delta_j} \right) > 0 \quad j = 1, 2 \dots m \quad k = 1, 2 \dots g$$

$$|\arg z_1| < \frac{1}{2} \pi \Delta_1 \quad |\arg z_2| < \frac{1}{2} \pi \Delta_2 \quad \Delta_1 \Delta_2 > 0$$

$$\Delta_1 = \sum_1^{M_1} b_j - \sum_{M_1+1}^{Q_1} b_j + \sum_1^{N_1} a_j - \sum_{N_1+1}^{P_1} a_j$$

$$\Delta_2 = \sum_1^{M_2} d_j - \sum_{M_2+1}^{Q_2} d_j + \sum_1^{N_2} c_j - \sum_{N_2+1}^{P_2} c_j$$

**Result III**

$$\begin{aligned} & \mathcal{L} \left\{ e^{-nt} t^h H_{P,Q}^{M,N} \left[ zt^k / \begin{matrix} (a_j, \alpha_j)_{1,P} \\ (b_j, \beta_j)_{1,Q} \end{matrix} \right], P \right\} \\ &= (P+a)^{-1-h} H_{P+1,Q}^{M,N+1} \left[ z(P+a)^{-k} / \begin{matrix} (-h, k) (a_j, \alpha_j)_{1,P} \\ (b_j, \beta_j)_{1,Q} \end{matrix} \right] \end{aligned}$$

Provided  $\operatorname{Re}(p) > 0$   $1 \geq \lambda > a$  and  $\operatorname{Re}(1+\alpha) > 0$

**Proof I** First Taking by mellin barnes type contour integral for H- function for one variable and then convolution of laplace transform for H-function and get required result.

**Proof II** Taking by mellin barnes type contour integral for H- function for two variables and then using beta function we get required result.

**Proof III** same as proof I

**3. Conclusion**

From this Paper we get some many solution of integral equation of convolution from having H – Function of one or more variables

**4. Acknowledgement**

The author is highly grateful to Dr. Atul Garg for their valuable help and suggestions to improve this paper

**5. Refrence**

1. Saxena R.K. , and Kumbhat R.K. , Integral operators H- Function, Indian J . Pure Appl. Maths 5(1974). 1-6
2. Srivasatava H.M. , Gupta K.C. , and Goyal S.P. The H- Function of one and two variables with applications south Asian Publishers. New Dehli and Madras (1992)
3. Erdely A. etal. Tabres of Integral Transforms, Mc Graw Hill- Book Company , New York , Vol. I. (1954).
4. Melachlan , N.W. : laplace Transform and their application to Differential equations, Dover, Publications New York (1962)
5. Muhammad : Special function and its Application, Ph.D. Thesis, University of Calcuta ,(1987)
6. kumbhat R.K. and K.M. khan Arif : Convolution of an Integral equation with the I-Function as its Kernel J.indian Acad. Math vol.23 No. 2 (2001)